



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

II. Solution by J. SCHEFFER, A. M., Hagerstown, Md., and A. H. HOLMES, Brunswick, Me.

By clearing of fractions, adding the four resulting equations and solving for $x-a$, we have

$$x-a=b(\lambda-\mu-\nu+s)/(\lambda+\mu+\nu+s)\dots(1).$$

Also, after the equations are cleared of fractions, by subtracting the sum of the second and third from the sum of the first and fourth, and solving for $x+a$, we have

$$x+a=b(\lambda+\mu+\nu+s)/(\lambda-\nu-\mu+s)\dots(2).$$

Multiplying (1) by (2), we have $x^2-a^2=b^2$ or $x^2=a^2+b^2$.

Also solved by G. W. Greenwood and G. B. M. Zerr.

Professor Greenwood furnished two solutions, the first of which made use of the vanishing of the determinant of the coefficients of λ, μ, ν, s in order that the four equations be consistent. The second solution was obtained by finding the values of $\mu/\nu, \nu/\lambda, s/\lambda$ in any three of the four equations and substituting them in the remaining equation.

Dr. Zerr solved each of the four equations for $x\lambda, x\mu, x\nu$, and xs , respectively, and added the square of the resulting equations. By dividing this last equation by $x^2\lambda^2, x^2$ is immediately found.

Professors Zerr and Greenwood also sent in solutions of 271.

GEOMETRY.

300. Proposed by J. J. QUINN, Ph. D., Scottdale, Pa.

Trisect an angle by means of a tractrix.

No solution has been received.

301. Proposed by PROF. R. D. CARMICHAEL, Anniston, Ala.

Apply the locus of $r=a(1+2\cos\theta)$ to the trisection of an angle. Describe the curve by continuous motion.

Solution by B. F. FINKEL, Ph. D., Drury College, Springfield, Mo.

To describe the curve by continuous motion, we may construct an instrument of the following description:

Take a bar AP and cut a path, indicated in the figure by the continuous white line, equal in length to four times the radius OA of the circle. Fasten the radius OQ at the middle point, Q , of the bar AP by means of a joint. Fasten the other end, O , of the radius at O . At P , a distance from Q equal to OQ , insert a pen point. Then if the path in the bar is set over a fixed pin at A , at a distance from O equal to OQ , the point P will describe the curve as the bar is made to move so as to keep A in the path of the bar.

